

Binary trees, coproducts, and integrable systems

B Erbe¹ and H J Schmidt²

¹Department of Physics, University of Regensburg, Regensburg, Germany

E-mail: bjoern.erbe@physik.uni-regensburg.de

²Department of Physics, University of Osnabrueck, Osnabrueck, Germany

E-mail: hschmidt@uos.de

Abstract. We provide a unified framework for the treatment of special integrable systems which we propose to call “generalized mean field systems”. Thereby previous results on integrable classical and quantum systems are generalized. Following Ballesteros and Ragnisco, the framework consists of a unital algebra with brackets, a Casimir element, and a coproduct which can be lifted to higher tensor products. The coupling scheme of the iterated tensor product is encoded in a binary tree. The theory is exemplified by the case of a spin octahedron. The relation to other generalizations of the coalgebra approach is discussed.

PACS numbers: 02.30.Ik, 75.10.Jm 20.00, 75.10.Hk

1. Introduction

In classical mechanics “complete integrability” can be precisely defined in terms of the Arnol’d-Liouville theorem [1]. The corresponding generalization of this concept to quantum theory has not yet been achieved. Nevertheless, there exists a rich literature on integrable quantum systems under various headlines such as Yang-Baxter equations [2], algebraic Bethe ansatz [3] and quantum groups [4]. Aside from this mainstream of research there are different theories of integrable systems which could be characterized as “generalized mean field systems” (GMFS) [5], [6]. It is the aim of the present article to provide a general framework for the description of such systems.

The prototype of the systems in question is a spin system where the spins are (Heisenberg-) coupled to each other with equal strength. It turns out that each spin will move exactly as if it would be under the influence of a uniform magnetic field. This justifies the above characterization as (generalized) “mean field systems”. The first generalizing step would be to consider systems which consist of uniformly coupled integrable subsystems. This property can be recursively applied. The underlying sequence of partial uniform couplings is most conveniently encoded in a binary tree, the leaves of which correspond to the smallest constituents of the system, see [6]. For example, the uniform coupling of three pairs of spins can be described by the tree of figure 1 and gives rise to an integrable spin octahedron, see figure 5.

Another generalization into a different direction is based upon the insight that at the core a GMFS consists of a unital algebra \mathcal{A} with a bracket $[\cdot, \cdot]$ and a co-multiplication Δ , which can be lifted to tensor products of \mathcal{A} and applied to a Casimir (central element) $c \in \mathcal{A}$, see [5]. One then considers representations of \mathcal{A} generated by certain commutation relations where the bracket $[\cdot, \cdot]$ will either be represented by a Poisson bracket (classical case) or by the commutator of matrices (quantum case).

In our paper we simplify and generalize the approaches of [5] and [6]. Thereby the restriction to Heisenberg spin systems in [6] is abolished by incorporating the coalgebra ansatz of [5]. Vice versa, the theory of [5] will be reformulated by using the language of binary trees, and generalized from “homogeneous trees” to general ones. We also found that the postulate in [5] of Δ being “co-associative” is superfluous, but see Appendix A. After the first publication of the coalgebra approach [5] various generalizations have been proposed, see [7], [8], [9], [10] and [11]. We will comment on the relation of our approach to these generalization at the appropriate places in the paper and in two appendices. The obvious generalization of assuming several Casimir elements instead of a single one will be neglected here.

The paper is organized as follows. In section 2 we collect some definitions concerning binary trees which are needed later. Section 3 is devoted to the algebraic prerequisites




Figure 1. Example of a binary tree with root r , six leaves ℓ_1, \dots, ℓ_6 and further nodes $\alpha, \beta, \gamma, \delta$.

including the coproduct Δ and its lift Δ^T to higher tensor products given by a binary tree T . In section 4 we apply these tools to the theory of integrable systems and prove the main result, theorem 1, which is analogous to prop. 1 of [5] and provides a number of commuting observables which is in many examples sufficient to guarantee complete integrability. In section 5 we discuss the elementary example of a Heisenberg spin octahedron in order to illustrate the application of the abstract theory. Some remarks of the corresponding Gaudin spin system and on the connection to other approaches follow. Two appendices on the issues of superintegrability and the recent loop coproduct approach close the paper.

2. Trees

We consider finite, binary trees T , shortly called “trees”. Recall that these consist of a set of “nodes” $\mathcal{N}(T)$, such that all nodes $n \in \mathcal{N}(T)$, except the “leaves” $\ell \in \mathcal{L}(T)$, are connected to exactly two “children” $c_1(n), c_2(n)$, mixing the metaphors of horticulture and genealogy. We have to distinguish between the “left child” c_1 and the “right child” c_2 . Due to this distinction, the leaves of a binary tree can be arranged in a natural order from left to right and hence be labelled by “ ℓ_1 ” to “ ℓ_L ”. All nodes, except the “root” $r(T)$, are children of other nodes. By definition, different nodes have different children, see figure 1. As a tree, T is a connected graph without cycles.

The simplest tree \bullet consists of only one root. The next simplest one \mathbb{V} has three nodes, that is, one root and two leaves. If T_1 and T_2 are (disjoint) trees, then $\mathbb{V}(T_1, T_2)$




Figure 2. Union $V(T_1, T_2)$ of two binary trees T_1 and T_2 .




Figure 3. Examples of “homogeneous” binary trees.

will be the tree obtained by identifying the leaves of V with the roots $r(T_1)$ and $r(T_2)$, see figure 2.

Obviously, each tree can be obtained from copies of \bullet by recursively applying the operation $V(T_1, T_2)$. This opens the possibility to provide recursive definitions and proofs in the theory of trees. A tree T will be called “homogeneous” if it is of the form

$$T = V(\dots(V(V(\bullet, \bullet), \bullet), \dots), \bullet) \tag{1}$$

$$\text{or } T = V(\bullet, V(\bullet, \dots, V(\bullet, \bullet) \dots)) , \tag{2}$$

see, for example, figure 3. The tree of figure 1 is not homogeneous.

Binary trees are used in various parts of physics, e. g. in chaos theory [12], computational physics [13], or in the theory of spin networks [14]. Here we utilize these structures for encoding the coupling schemes of certain integrable spin systems, similarly as in [15], [16] and [6].




Figure 4. Binary tree $T = T_1 \circ_{\ell} T_2$ obtained by grafting T_2 on T_1 .

The following lemma can be easily proved.

Lemma 1 $N(T) \equiv |\mathcal{N}(T)| = 2|\mathcal{L}(T)| - 1 \equiv 2L(T) - 1$.

A sub-tree $S \subset T$ is given by a subset of nodes of T , which, according to their connections inherited from T again form a tree. For example, if $n \in \mathcal{N}(T)$ then $T(n)$ will denote the maximal sub-tree of T with the root n . Let $\mathcal{L}(n) \equiv \mathcal{L}(T(n))$. If $n, m \in \mathcal{N}(T)$, then either $\mathcal{L}(n) \subset \mathcal{L}(m)$ or $\mathcal{L}(m) \subset \mathcal{L}(n)$, or $\mathcal{L}(n) \cap \mathcal{L}(m) = \emptyset$. In the former two cases m and n will be called “connected”, in the latter case “disjoint”.

If T_1 and T_2 are (disjoint) trees and $\ell \in \mathcal{L}(T_1)$, then $T = T_1 \circ_{\ell} T_2$ will denote the tree obtained by “grafting”, i. e. by identifying the root $r(T_2)$ with the leaf ℓ of T_1 , see figure 4.

3. Coproducts

In this paper we often will consider the classical and the quantum case simultaneously. In both cases, the physical observables are obtained by suitable representations of an abstract unital algebra (\mathcal{A}, e) and its tensor products. In the quantum case, \mathcal{A} will be an associative, non-abelian algebra with commutator $[a, b] = ab - ba$, $a, b \in \mathcal{A}$, and its physical representation is given in terms of finite-dimensional matrices. Typical examples are cases where \mathcal{A} is defined as the universal enveloping algebra of some semi-simple Lie algebra. In the classical case, \mathcal{A} will be an abelian algebra together with an

abstract Poisson bracket $\{, \}$, see [5]. Representations of \mathcal{A} are then given by the algebra of smooth functions of some phase space together with the usual Poisson bracket.

To cover both cases, the commutator/Poisson bracket will be denoted by $[a, b]$, $a, b \in \mathcal{A}$. It makes $(\mathcal{A}, [,])$ into a Lie algebra and will act as an derivation on the associative product on \mathcal{A} . We will always consider algebras endowed with a bracket of one of these two kinds and the corresponding homomorphisms, that is, linear algebra homomorphism w. r. t. both multiplications.

If $\mathcal{A}_1, \mathcal{A}_2$ are two algebras as explained above, then $\mathcal{A}_1 \otimes \mathcal{A}_2$ will denote the algebraic tensor product, physically describing a composite system. It will be again a unital algebra with brackets upon linearly extending the definitions

$$(a \otimes b)(c \otimes d) = (ac) \otimes (bd) \quad (3)$$

and

$$[(a \otimes b), (c \otimes d)] = [a, c] \otimes \frac{bd + db}{2} + \frac{ac + ca}{2} \otimes [b, d]. \quad (4)$$

If $A : \mathcal{A}_1 \rightarrow \mathcal{A}_1$ and $B : \mathcal{A}_2 \rightarrow \mathcal{A}_2$ are morphisms as explained above, then also $A \otimes B : \mathcal{A}_1 \otimes \mathcal{A}_2 \rightarrow \mathcal{A}_1 \otimes \mathcal{A}_2$ will be such a morphism.

The ‘‘coproduct’’ Δ will be a morphism

$$\Delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}, \quad (5)$$

that is, a linear algebra morphism plus a Poisson bracket morphism in the classical case. Usually, a coproduct is additionally required to be ‘‘co-associative’’, see [5], but this property will not be needed in the main part of the present paper, hence here we use the term ‘‘coproduct’’ in a more general sense. In Appendix A, co-associativity and co-commutativity of Δ will be assumed to extend the set of integrals of motion (super-integrability).

The crucial construction for integrability, as considered here, is the lift of the coproduct to higher order tensor products given by a tree T . To this end we first define \mathcal{A}^T recursively by

$$\mathcal{A}^\bullet = \mathcal{A} \quad (6)$$

$$\mathcal{A}^{\vee(T_1, T_2)} = \mathcal{A}^{T_1} \otimes \mathcal{A}^{T_2}. \quad (7)$$

Sometimes it will be convenient to use the identification

$$\mathcal{A}^{T_1} = \mathcal{A}^{T_2} = \underbrace{\mathcal{A} \otimes \dots \otimes \mathcal{A}}_{L \text{ times}}, \quad (8)$$

if $L(T_1) = L(T_2) \equiv L$. W. r. t. this identification the canonical embedding

$$j_n : \mathcal{A}^{T(n)} \rightarrow \mathcal{A}^T, \quad n \in \mathcal{N}(T) \quad (9)$$

can be defined by

$$j_n(a) = e \otimes \dots \otimes a \otimes \dots \otimes e, \quad a \in \mathcal{A}^{T(n)}. \quad (10)$$

In the next step we define the lift of the coproduct $\Delta^T : \mathcal{A} \longrightarrow \mathcal{A}^T$ recursively by

$$\Delta^\bullet = \text{id}_{\mathcal{A}} \quad (11)$$

$$\Delta^{\vee(T_1, T_2)} = (\Delta^{T_1} \otimes \Delta^{T_2}) \circ \Delta, \quad (12)$$

and conclude:

Lemma 2 $\Delta^T : \mathcal{A} \longrightarrow \mathcal{A}^T$ is a (Poisson) algebra morphism.

Proof. By induction over T . The claim follows since the tensor product and the composition of (Poisson) algebra morphisms is again a (Poisson) algebra morphism. \square

Before formulating the main result we still need another definition. Let $n \in \mathcal{N}(T)$, then

$$\Delta^n \equiv j_n \circ \Delta^{T(n)} : \mathcal{A} \longrightarrow \mathcal{A}^T. \quad (13)$$

We note that the generalization of the coalgebra approach to comodule algebras [7] where Δ^n is replaced by a suitable map $\mathcal{A} \longrightarrow \mathcal{A} \otimes \mathcal{B} \otimes \dots \otimes \mathcal{B}$ is only possible for homogeneous trees.

4. Integrable systems

Remember the grafting of trees $T = T_1 \circ_\ell T_2$ explained in section 2. It gives rise to a corresponding composition of lifted coproducts in the following sense:

Lemma 3 Let $T = T_1 \circ_\ell T_2$, $x \in \mathcal{A}$ and write $\Delta^{T_1}(x) = \sum_i x_{i, \ell_1} \otimes \dots \otimes x_{i, \ell_\mu} \otimes \dots \otimes x_{i, \ell_{L_1}}$, where $\ell_1, \dots, \ell_{L_1}$ denote the leaves of T_1 and $\ell_\mu = \ell$. Then $\Delta^T(x) = \sum_i x_{i, \ell_1} \otimes \dots \otimes \Delta^{T_2}(x_{i, \ell_\mu}) \otimes \dots \otimes x_{i, \ell_{L_1}}$.

Proof. By induction over T_1 . \square

Further one can easily prove the following lemma:

Lemma 4 Let $n, m \in \mathcal{N}(T)$ be disjoint, i. e. $\mathcal{L}(n) \cap \mathcal{L}(m) = \emptyset$ and $x, y \in \mathcal{A}$. Then $[\Delta^n(x), \Delta^m(y)] = 0$.

Proof. $\Delta^n(x)$ and $\Delta^m(y)$ live in disjoint factors of the tensor product \mathcal{A}^T , since they are of the form $\Delta^n(x) = \sum_i e \otimes \dots \otimes_{\ell \in \mathcal{L}(n)} x_{i, \ell} \otimes \dots \otimes e$ and $\Delta^m(y) = \sum_j e \otimes \dots \otimes_{\ell \in \mathcal{L}(m)} y_{j, \ell} \otimes \dots \otimes e$. Hence they commute. \square

Now we are ready to formulate the main result.

Theorem 1 Let $n, m \in \mathcal{N}(T)$ such that $\mathcal{L}(m) \subset \mathcal{L}(n)$, $x \in \mathcal{A}$ and $c \in \mathcal{A}$ be a central element. Then $[\Delta^n(x), \Delta^m(c)] = 0$.

Proof. We introduce the canonical partial embedding $j_m^n : \mathcal{A}^{T(m)} \longrightarrow \mathcal{A}^{T(n)}$ such that $j_m = j_n \circ j_m^n$. It follows that

$$\begin{aligned} & [\Delta^n(x), \Delta^m(c)] \\ &= [j_n \circ \Delta^{T(n)}(x), j_m \circ \Delta^{T(m)}(c)] \end{aligned} \quad (14)$$

$$= j_n ([\Delta^{T(n)}(x), j_m^n \circ \Delta^{T(m)}(c)]) \quad (15)$$

and thus it suffices to show that

$$[\Delta^{T(n)}(x), j_m^n \circ \Delta^{T(m)}(c)] = 0. \quad (16)$$

By applying lemma 3 to the sub-tree $T(n)$ we write

$$T(n) = T_1 \circ_m T(m), \quad m = \ell_\mu \in \mathcal{L}(T_1) \quad (17)$$

and conclude

$$\begin{aligned} & \Delta^{T(n)}(x) \\ &= \sum_i x_{i,1} \otimes \dots \Delta^{T(m)}(x_{i,\mu}) \otimes \dots x_{i,L_1}. \end{aligned} \quad (18)$$

Hence

$$\begin{aligned} & [\Delta^{T(n)}(x), j_m^n \circ \Delta^{T(m)}(c)] \\ &= \sum_i x_{i,1} \otimes \dots [\Delta^{T(m)}(x_{i,\mu}), \Delta^{T(m)}(c)] \otimes \dots x_{i,L_1} \\ &= \sum_i x_{i,1} \otimes \dots \Delta^{T(m)}([x_{i,\mu}, c]) \otimes \dots x_{i,L_1} \\ &= 0, \end{aligned} \quad (19)$$

since $[x_{i,\mu}, c] = 0$. □

This theorem generalizes prop. 1 of [5] to arbitrary, not necessarily homogeneous trees. In order to guarantee complete integrability in the sense of the Arnol'd-Liouville theorem for $2L$ -dimensional phase spaces (which is satisfied for spin systems, see section 5) we would need L pairwise commuting observables (“integrals in involution”). These are provided by the $\Delta^n(c)$ for each node n which is not a leaf. By lemma 1, there are exactly $N(T) - L(T) = L(T) - 1$ such nodes. In [5] the remaining observable is chosen as the Hamiltonian H . In the context of quantum spin systems another choice would be more appropriate, namely $\Delta^r(x)$ with a suitable $x \in \mathcal{A}$. The Hamiltonian H could then be chosen as any element of the algebra generated by the $\Delta^n(c)$ and $\Delta^r(x)$.

In the general case the dimension of the phase space depends on the symplectic realization of $(\mathcal{A}, \{, \})$ and the choice of the symplectic leaves. A thorough discussion of these questions, which also applies to the binary tree approach, including issues of superintegrability can be found in [9].

5. Examples and Outlook

In order to explain the application of theorem 1 to integrability of quantum systems we consider the elementary example of a spin octahedron, figure 5, with Heisenberg Hamiltonian, following [6]. We chose as \mathcal{A} the universal enveloping algebra of the Lie algebra $su(2)$. More concretely: We consider three generators X_1, X_2, X_3 satisfying the abstract commutations relations

$$[X_j, X_k] = i \sum_{\ell=1}^3 \epsilon_{j k \ell} X_\ell, \quad (20)$$

where $\epsilon_{j k \ell}$ denotes the completely anti-symmetric Levi-Civita symbol. \mathcal{A} is the set of all finite polynomials X of the standard form

$$X = \sum_{klm} c_{klm} X_1^k X_2^l X_3^m. \quad (21)$$

The product XY of two such polynomials is brought into the standard form (21) by successively applying the commutation relations (20). The unit element in \mathcal{A} is $e = X_1^0 X_2^0 X_3^0$. It follows that $c \equiv X_1^2 + X_2^2 + X_3^2$ commutes with all $X \in \mathcal{A}$.

The coproduct Δ is defined on the generators by

$$\Delta(X_i) = e \otimes X_i + X_i \otimes e \quad (22)$$

and then extended to general elements of the form (21) by employing the property of Δ being an algebra homomorphism. Thus, for example,

$$\Delta(c) = \Delta(X_1^2 + X_2^2 + X_3^2) \quad (23)$$

$$= \Delta(X_1)^2 + \Delta(X_2)^2 + \Delta(X_3)^2 \quad (24)$$

$$= \sum_{i=1}^3 (e \otimes X_i + X_i \otimes e)^2 \quad (25)$$

$$= e \otimes c + c \otimes e + 2 \sum_{i=1}^3 X_i \otimes X_i. \quad (26)$$

We further choose T as the binary tree of figure 1 and obtain the corresponding various commutations relations of theorem 1 being valid in the 6-fold tensor product $\bigotimes_{i=1}^6 \mathcal{A}$ where one usually chooses $\Delta^r(x) = \Delta^r(X_3)$. Next we consider the well-known $(2s+1)$ -dimensional irreducible matrix representation of (20) and denote the representations of the generators X_i by \mathbf{S}_i (“spin operator components”). In the 6-fold tensor product we denote the single spin components by \mathbf{S}_i^μ , $\mu = 1, \dots, 6$. In this representation, $c = s(s+1)\mathbf{1}$ and all commutation relations of theorem 1 remain valid. Note that $\Delta(c)$ becomes $(\mathbf{S}^{(1)} + \mathbf{S}^{(2)})^2$, which is no longer a constant, analogously for higher tensor products $\Delta^n(c)$. This shows, by the way, why it is advantageous to work in an abstract setting and to consider concrete representations only after the coproduct is defined. Note further, that we could slightly generalize the example by considering




Figure 5. The octahedral spin graph corresponding to the integrable Heisenberg Hamiltonian (27). Its coupling scheme is encoded in the binary tree of figure 1 as explained in the text.

different s for each factor of the tensor product.

Let $(\mathcal{V}, \mathcal{E})$ be the octahedral spin graph of figure 5 with its set of 6 vertices \mathcal{V} and the set of 12 edges \mathcal{E} . The corresponding Heisenberg Hamiltonian H can be written in various ways:

$$H = 2J \sum_{(\mu, \nu) \in \mathcal{E}} \mathbf{S}^{(\mu)} \cdot \mathbf{S}^{(\nu)} \quad (27)$$

$$= J \left(\left(\sum_{\mu \in \mathcal{V}} \mathbf{S}^{(\mu)} \right)^2 - (\mathbf{S}^{(1)} + \mathbf{S}^{(2)})^2 - (\mathbf{S}^{(3)} + \mathbf{S}^{(4)})^2 - (\mathbf{S}^{(5)} + \mathbf{S}^{(6)})^2 \right) \quad (28)$$

$$= J (\Delta^r(c) - \Delta^\alpha(c) - \Delta^\beta(c) - \Delta^\gamma(c)) , \quad (29)$$

where the root r and the nodes α, β, γ refer to the binary tree of figure 1. J is some appropriate coupling constant. It is crucial that H can be written as a linear combination of commuting observables according to theorem 1. In this respect the octahedral Heisenberg Hamiltonian (27) is only the simplest case; for example, a Zeeman term proportional to $\Delta^r(\mathbf{S}_3)$ could be added without losing integrability. The eigenvalues and common eigenvectors of the system of commuting observables $\Delta^n(c)$, $n \in \mathcal{N}(T)$, result from the well-known rules of coupling angular momenta involving Clebsch-Gordan coefficients. An explicit formula for the eigenvalues and eigenvectors of H and arbitrary

binary trees has been given in [6]. The example of the spin octahedron clearly shows the physical meaning of the binary tree T on which theorem 1 depends: T encodes the coupling scheme of systems which are completely integrable due to their structure of uniformly coupled subsystems.

With exactly the same algebraic considerations and the same tree as above, theorem 1 provides us with another very interesting integrable model:

$$\begin{aligned} H &= A \left(\mathbf{S}^{(1)} \cdot \sum_{i=3}^6 \mathbf{S}^{(i)} + \mathbf{S}^{(2)} \cdot \sum_{i=3}^6 \mathbf{S}^{(i)} \right) \\ &\quad + 2(A + J)\mathbf{S}^{(1)} \cdot \mathbf{S}^{(2)} \\ &= A (\Delta^r(c) - \Delta^\delta(c)) + J\Delta^\alpha(c), \end{aligned} \tag{30}$$

The (Gaudin) Hamiltonian H describes a central spin system with two central spins of exchange $2(A + J)$, coupled homogenously to a bath of four spins. Such a system can serve for example as a simplified model for the hyperfine interaction in a double quantum dot, see [17].

Apart from this physical meaning, it is interesting from a formal point of view. Besides the approach presented in this paper, systems can be integrable in the sense of algebraic Bethe ansatz. According to the ground breaking work of V. Drinfeld [4], this is based on quasicocommutative bialgebras, which essentially means that there is an element $R \in \mathcal{A} \otimes \mathcal{A}$ with

$$(\tau \circ \Delta)(x) \cdot R = R \cdot \Delta(x) \tag{31}$$

for all $x \in \mathcal{A}$. τ denotes the switch operator defined by linearly extending $\tau(a \otimes b) = b \otimes a$ and Δ a usual coproduct.

As this algebraic structure is somewhat similar to the one presented in this paper, the question arises, whether there is a connection between systems integrable in either sense. The above system, in contrast to the central spin system with one central spin, is not integrable by means of algebraic Bethe ansatz. Hence adding a second central spin destroys the Bethe ansatz, whereas integrability in the sense of theorem 1 remains unaffected.

Recently, a framework for integrability using so-called “loop coproducts” has been proposed [10], [11] which contains different previous approaches to integrability as special cases. It is, however, confined to the classical case. Some remarks on the relation between this approach and the present paper are included in the appendix B.

Appendix A: Superintegrability

The approach [5] to integrability via coalgebras has subsequently be extended to “super-integrability” [8], [9]. This roughly means that one is seeking for additional integrals of motion which, however, do not longer commute with the old ones. “Additional” means, in the classical case, that the new integrals of motion are functionally independent of the old ones. Typically, this functional independence cannot be shown in the general

setting, but only in concrete examples, see [8], [9]. Also in these references the role of co-associativity of Δ in connection to superintegrability has been stressed.

The question arises, whether these ideas can be transferred to the more general situation where the binary trees are not necessarily of homogeneous type. To this end we slightly refine the binary tree construct, following, for example, [15]. Recall that, due to the distinction between “left child” and “right child”, the leaves of a binary tree can be arranged in a natural order from left to right and hence be labeled by “ ℓ_1 ” to “ ℓ_L ”. Now assume that this labeling can be arbitrarily permuted. We will call the resulting structure a coupling tree. For example, the only binary tree with three nodes, \mathbf{V} , gives rise to two different coupling trees, denoted by $\mathbf{V}(\ell_1, \ell_2)$ and $\mathbf{V}(\ell_2, \ell_1)$. Generally, a coupling tree \tilde{T} can be represented as a pair $\tilde{T} = (T, \pi)$, where T is a binary tree with L leaves and $\pi \in \mathcal{S}_L$, a permutation of L elements. A coupling tree with L leaves can alternatively be construed as a monomic expression in the abstract variables ℓ_1, \dots, ℓ_L , such that each variable occurs exactly once. For example, the tree of figure 1, conceived as a coupling tree, corresponds to the expression $(\ell_1 \ell_2)((\ell_3 \ell_4)(\ell_5 \ell_6))$.

Most definitions and propositions of the sections 2 and 3 can be taken over directly or with minor modifications. It will be appropriate to reserve the union and grafting operations, see figures 2 and 4, to binary trees, and to obtain the corresponding coupling trees by adducing a suitable permutation of the leaves, as explained above. Note, that the definition of \mathcal{A}^T remains unchanged. We will extend the definitions (6) and (7) to coupling trees by

$$\Delta^{\tilde{T}} = \tilde{\pi} \circ \Delta^T, \quad (32)$$

where $\tilde{T} = (T, \pi)$ and $\tilde{\pi} : \mathcal{A}^T \longrightarrow \mathcal{A}^T$ denotes the natural representation of π by a permutation of factors of the tensor product. Following [15] we consider two operations on coupling trees, namely

$$\text{exchange: } \mathbf{V}(\ell_1, \ell_2) \rightleftharpoons \mathbf{V}(\ell_2, \ell_1) \quad (33)$$

$$\text{flop: } \mathbf{V}(\mathbf{V}(\ell_1, \ell_2), \ell_3) \rightleftharpoons \mathbf{V}(\ell_1, \mathbf{V}(\ell_2, \ell_3)), \quad (34)$$

see figure 6. We have the following

Proposition 1 *Let T_1 and T_2 be two coupling trees with $\mathcal{L}(T_1) = \mathcal{L}(T_2)$. Then T_1 can be transformed into T_2 by a finite sequence of exchanges and flops operating on subtrees.*

We will skip the proof which is lengthy but straight forward. Note that, in the language of monomials, the proposition says that any two monomials with the variables ℓ_1, \dots, ℓ_L occurring exactly once can be transformed into each other by applying the rules of commutativity and associativity of the multiplication.

We will say that the coproduct $\Delta : \mathcal{A} \longrightarrow \mathcal{A} \otimes \mathcal{A}$ is co-commutative iff $\Delta^{\mathbf{V}(\ell_1, \ell_2)} = \Delta^{\mathbf{V}(\ell_2, \ell_1)}$ and co-associative iff $\Delta^{\mathbf{V}(\mathbf{V}(\ell_1, \ell_2), \ell_3)} = \Delta^{\mathbf{V}(\ell_1, \mathbf{V}(\ell_2, \ell_3))}$. Note that the




Figure 6. Illustration of the operations “exchange” and “flop” on coupling trees.

coproduct defined in (22) is co-commutative as well as co-associative. Obviously, Δ is co-commutative and co-associative iff Δ^T is invariant under exchanges and flops operating on sub-trees of T . Together with proposition 1 we obtain:

Proposition 2 *If Δ is co-commutative and co-associative and $\mathcal{L}(T_1) = \mathcal{L}(T_2)$ then $\Delta^{T_1} = \Delta^{T_2}$.*

Moreover, if $n_1 \in \mathcal{N}(T_1)$ and $n_2 \in \mathcal{N}(T_2)$ such that $\mathcal{L}(n_1) = \mathcal{L}(n_2)$ then $\Delta^{n_1} = \Delta^{n_2}$.

Now let $\mathcal{L}(T_1) = \mathcal{L}(T_2)$ and consider the involutive sub-algebra $\mathcal{C}_1 \subset \mathcal{A}^{T_1}$ generated by the elements $\Delta^n(c)$, $n \in \mathcal{N}(T_1)$ and $\Delta^{r_1}(x)$, analogously for $\mathcal{C}_2 \subset \mathcal{A}^{T_2}$, see section 4. Define

$$CN(T_1, T_2) = \{(n_1, n_2) | n_1 \in \mathcal{N}(T_1), n_2 \in \mathcal{N}(T_2), \mathcal{L}(n_1) = \mathcal{L}(n_2)\}, \quad (35)$$

and \mathcal{C}_{12} as the sub-algebra of $\mathcal{C}_1 \cap \mathcal{C}_2$ generated by the elements $\Delta^{n_1}(c)$, $\Delta^{r_1}(x)$ or, equivalently, by the $\Delta^{n_2}(c)$, $\Delta^{r_2}(x)$, where (n_1, n_2) runs through $CN(T_1, T_2)$. Then we conclude the main result of this appendix:

Theorem 2 *Let Δ be co-commutative and co-associative and $H \in \mathcal{C}_{12}$, then $[H, K] = 0$ for all $K \in \mathcal{C}_1 \cup \mathcal{C}_2$.*

The scenario for superintegrability considered in [8], [9] results as a special case of theorem 2 in the following sense. Let T_1 be the “left-homogeneous tree” and T_2 the “right-homogeneous tree” represented in figure 3. Then $CN(T_1, T_2) = \{(r_1, r_2)\}$ and \mathcal{C}_{12} is the algebra generated by $\Delta^{r_1}(x) = \Delta^{r_2}(x)$. Note that in this case T_1 can be transformed into T_2 using only flops operating on sub-trees, hence the assumption of Δ being co-commutative will be superfluous.

Appendix B: Loop coproducts

Recently, a framework for integrability using so-called “loop coproducts” has been proposed by F. Musso [10], [11] which contains different previous approaches to integrability as special cases, namely the coalgebra approach [5], the linear r-matrix formulation and formulations using Sklyanin or reflection algebras. It is, however, confined to the classical case. Nevertheless, one may ask whether, in the classical case, the loop coproduct approach also includes the generalization of the coalgebra approach we have given in this paper.

At first glance, the answer seems to be “no”, since the corresponding derivation in [10] of the coalgebra as a special case utilizes the co-associativity of Δ , which is not needed in our theorem 1. Here we neglect the differences due to the assumption in [10] and [11] that the algebra \mathcal{A} has a finite number of generators. A closer inspection, however, reveals that co-associativity is not necessary.

The loop coproduct approach [11] is based on a family of maps $\Delta^{(k)} : \mathcal{A} \longrightarrow \mathcal{B}$, $k = 1, \dots, m$ and postulates different properties of these maps for the cases $i < k$ (or $k < i$) and $i = k$. For comparison we have to set $\mathcal{B} = \mathcal{A}^T$. In our approach, the set of nodes is only partially ordered by the definition $i \prec k$ iff $\mathcal{L}(i) \subset \mathcal{L}(k)$. However, \prec can be extended to a linear order $<$ such that $i < k$ implies $\mathcal{L}(i) \subset \mathcal{L}(k)$. For $i < k$ we have either $\mathcal{L}(i) \cap \mathcal{L}(k) = \emptyset$ or $\mathcal{L}(i) \subset \mathcal{L}(k)$. In the first case $[\Delta^i(x), \Delta^k(y)] = 0$ for all $x \in \mathcal{A}$ due to lemma 4. In the second case lemma 3 implies

$$[\Delta^i(x), \Delta^k(y)] = \sum_j f_j [\Delta^i(x_j), \Delta^i(y)] \quad (36)$$

for all $x, y \in \mathcal{A}$ and some suitable $f_j \in \mathcal{B}$, $x_j \in \mathcal{A}$. Hence, in both cases condition (4) of [11] is satisfied. If $i = k$, condition (5) of [11] follows since Δ^i is a (Poisson) algebra homomorphism in our theory.

We conclude that, in the case of classical mechanics and up to minor differences in the formulations, the loop coproduct theory [10, 11] contains the binary tree approach as a special case. Nevertheless, the binary tree approach has, to our opinion, its virtues as a constructive method particularly adapted to quantum spin systems.

Acknowledgements

We are indebted to a referee for hints to the literature concerning generalizations of [5] and to Roman Schnalle for references on coupling trees.

References

- [1] V. I. Arnol'd. *Mathematical Methods of Classical Mechanics*. Springer-Verlag, Berlin, 1978.
- [2] R. Baxter. *Exactly Solved Models in Statistical Mechanics*. Academic Press, London – New York, 1982.
- [3] L.D. Faddeev. How algebraic Bethe Ansatz works for integrable model. *Les Houches lectures*, hep-th/9605187, 1995.
- [4] V. G. Drinfeld. Quantum Groups. in: A. M. Gleason (ed.), *Proceedings of the International Congress of Mathematicians; Berkeley, California, USA, 1986*, 798-820, 1987.
- [5] A. Ballesteros and O. Ragnisco. A systematic construction of completely integrable hamiltonians from coalgebras. *J. Phys. A* 31:3791-3813, 1998.
- [6] R. Steinigeweg and H.-J. Schmidt. Heisenberg-integrable spin systems. *Math. Phys. Analysis and Geom.* 12, 1, 19-45, 2009.
- [7] A. Ballesteros, F. Musso, and O. Ragnisco. Comodule algebras and integrable systems. *J. Phys. A* 41:304028, 2008.
- [8] A. Ballesteros, F. J. Herranz, F. Musso, and O. Ragnisco. Superintegrable deformations of the Smorodinsky-Winternitz Hamiltonian. In: P. Tempesta et al (eds.) *Superintegrability in Classical and Quantum Systems*, CRM Proceedings & Lecture Notes, vol. 37, AMS, 2004.
- [9] A. Ballesteros and A. Blasco. N -dimensional superintegrable systems from symplectic realizations of Lie coalgebras. *J. Phys. A* 41:304028, 2008.
- [10] F. Musso. Loop coproducts. Preprint nlinSI/0907.4927, 2009.
- [11] F. Musso. Generalization of the r-matrix formulation through loop coproducts. Preprint nlinSI/0910.1047, 2009.
- [12] J. A. Ketoja and J. Kurkijrvi. Binary tree approach to scaling in unimodal maps *J. Stat. Phys.* 75, 3-4, 643-668, 2009
- [13] J. Waltz, G. L. Page, S. D. Milder, J. Wallin, and A. Antuns. A performance comparison of tree data structures for N-body simulation. *J. Comp. Phys.* 178, 1, 1-14, 2002
- [14] T. Tufarelli and V. Giovannetti. High fidelity state transfer in binary tree spin networks *Phys. Rev. A*, 79, 2, 022313, 2009
- [15] V. Fack, S. N. Pitre, and J. Van der Jeugt. New efficient programs to calculate general recoupling coefficients. Part I: Generation of a summation formula. *Comp. Phys. Comm.* 83:275-292, 1994.
- [16] R. Schnalle and J. Schnack. Numerically exact and approximate determination of energy eigenvalues for antiferromagnetic molecules using irreducible tensor operators and general point-group symmetries *Phys. Rev. B*, 79, 104419, 2009
- [17] J. Schliemann, A. Khaetskii and D. Loss. Electron spin dynamics in quantum dots and related nanostructures due to hyperfine interaction with nuclei. *J. Phys. : Condens. Matter* 15:1809-1833, 2003.